

MEDfacts

An Educational Health Series From National Jewish Health®



SARCOIDOSIS

Sarcoidosis is a chronic disease that can affect any organ in the body, but most commonly affects the lungs. Very small (microscopic) clusters of inflammation or white cells, called granulomas, are seen in the organs affected with sarcoidosis. These granulomas may clear up on their own, or may cause organ dysfunction and scarring. While there is no cure for sarcoidosis at this time, the disease can be managed to minimize organ dysfunction.

Sarcoidosis is likely to have more than one cause. Sarcoidosis is most common in young people between the ages of 20 and 40. However, it can affect any age group and race. About 10 to 40 of 100,000 people develop sarcoidosis. Sarcoidosis is not contagious.

What Are the Signs and Symptoms?

Sarcoidosis symptoms can vary greatly, and up to half of people with sarcoidosis have no symptoms when the illness is diagnosed. A person with sarcoidosis may have:

- No symptoms (asymptomatic)
- Only vague symptoms of a general nature, such as weight loss, night sweats, fatigue and low grade fever. High fever is less common, but can still occur.
- Symptoms caused by the affected organ.

The lungs are the most common organ affected by sarcoidosis. However, any organ can be affected. More than one organ can be involved. Signs and symptoms associated with specific organ involvement can include:

Lungs

Inflammation in the lungs can cause shortness of breath, wheezing or cough (often a dry cough). In some people, the symptoms go away. In others, there can be permanent scarring and persistent symptoms.

Lymph Nodes

Enlargement of various lymph nodes can occur, especially the

REGISTER NOW

to Receive **FREE** Health
Tips, Research Findings,
Ways to Support Our
Mission, News &
More in Your E-Mail Box

Visit njhealth.org/e-news
for more information.

Follow us online!

facebook

facebook.com/NJHealth

YouTube

youtube.com/NationalJewish

twitter

twitter.com/njhealth

lymph nodes in the chest, although these may not cause symptoms. Sometimes, lymph nodes can be associated with pain.

Eye

Inflammation of the eye can lead to redness, pain, dry eyes and sensitivity to light. Blurred vision also can occur. In some cases there can be eye involvement with no obvious visual problems. It is important that an eye doctor perform an eye exam regularly to determine if there is eye involvement.

Skin

Skin may appear as raised, pink or purplish areas or as painful nodules under the skin.

Bone

Bone involvement is often detected incidentally. Sometimes it can cause pain and rarely fractures.

Spleen and Liver

Enlargement of the spleen or liver that a doctor can feel during a physical exam can occur. The only abnormality may be seen on liver blood tests.

Heart

Heart involvement is thought to occur in up to 25 – 40 percent of people with sarcoidosis. It can be difficult to diagnose. Heart involvement can occur without symptoms. It can also appear with heart rhythm abnormalities (too fast or too slow). These can cause heart palpitations and lightheadedness. Heart involvement can also lead to problems with the heart muscle's ability to pump blood or heart failure.

Brain and Nervous System

Granulomas can develop in the brain and the nerves and cause many symptoms. Symptoms may include loss of sensation, loss of muscle strength, headaches and dizziness. This affects about one in 100 people with sarcoidosis.

Salivary Gland

The salivary gland can be involved. People with salivary involvement may have trouble with a dry mouth.

Kidneys and Calcium

Kidney involvement is not common. It can cause kidney damage. In addition, calcium levels may be elevated in the urine, leading to kidney stones. High calcium levels in the blood can cause constipation and generalized weakness.

How Is Sarcoidosis Diagnosed?

The first step in diagnosing sarcoidosis is a thorough evaluation. The inflamed microscopic granulomas seen in the affected organ with sarcoidosis are similar to those in other diseases such as tuberculosis, fungal diseases, berylliosis and farmer's lung. Because of this, a careful evaluation is important to rule out other diagnoses that can look like sarcoidosis. Only after the known causes of granulomas have been ruled out is the diagnosis of sarcoidosis made.

An evaluation to detect sarcoidosis should include the following:

Thorough Medical Examination: This can help rule out other diseases that may be similar to sarcoidosis.

Chest X-ray: Doctors look at chest X-rays for evidence of enlarged lymph nodes and small round spots in the lung caused by the clusters of inflammation. "Staging" can help

the doctor determine the degree of lung involvement in sarcoidosis. A scale of 0 – 4 is commonly used, with 4 having the highest amount of lung involvement.

Pulmonary Function Tests: These breathing tests give an indication of the severity of lung disease. There is nothing unique about sarcoidosis on these tests, so they do not substitute for other, more specific tests. Pulmonary function tests can show obstruction of airflow out of the lungs, restriction of the lung's ability to take in air and a decrease in the transport of oxygen from the lung into the bloodstream. The most important types of breathing tests in sarcoidosis are spirometry, lung volumes and diffusing capacity. In some cases, measurement of blood oxygen levels during an exercise test is done.

Tissue Biopsy: A microscopic exam of tissue samples from the lungs or other affected organs is also needed to be sure of the diagnosis and to exclude other causes. A bronchoscopy can obtain this tissue. A bronchoscopy is a procedure in which the doctor places a narrow tube through the nose and into the airways. Sometimes the diagnosis is made by obtaining tissue samples from the skin, liver or enlarged lymph nodes.

Bronchoalveolar Lavage: When a bronchoscopy is done, a small part of lung can be washed (lavaged) in order to obtain some cells of the immune system from the lung. By counting the types of cells in lavage fluid, it is possible to get an estimate of how inflamed the lungs are and whether the type of inflammation is characteristic of sarcoidosis.

Eye Exam: A slit lamp exam by an eye doctor (ophthalmologist) is an important part of an eye exam to detect inflammation.

CT (Computerized Tomography) Scan: A CT scan is a detailed type of X-ray. The CT scan may make it possible to see lymph nodes and scars in the lung when regular chest X-rays sometimes cannot.

Calcium Levels in the Blood and Urine: Regulation of calcium may be disturbed in sarcoidosis. This results in too much calcium in the blood and/or urine. Exposure to the sun, calcium and vitamin D supplements and high dairy intake can stimulate this process. A 24-hour urine sample can measure the calcium level in the urine. A blood sample can be drawn for levels of calcium in the blood. High calcium levels in the blood can be seen with symptoms that may include fatigue, abdominal pain/constipation and mental foginess. High calcium levels can also lead to kidney dysfunction.

PET (Positron Emission Tomography) Scan: A radioactive labeled sugar is given intravenously. The person is placed in a special scanning machine to see where the radioactive labeled sugar accumulated. Areas of active inflammation take up the radioactive labeled sugar and are detected with the scanning machine. This gives your doctor a better idea of which areas or organs might be involved with sarcoidosis.

Heart Testing: The doctor may order an electrocardiogram (EKG) to evaluate the electrical system of the heart. An echocardiogram may also be done to evaluate the structure and function of the heart. Sometimes, the doctor might order further tests for the heart if there is suspicion of heart involvement with sarcoidosis. These tests might include a Holter monitor and/or a heart MRI (magnetic resonance imaging).

What Is the Treatment?

Up to one-half of the people diagnosed with sarcoidosis improve without treatment. Those who do not improve are often placed on medicine to reduce inflammation. Many people will recover, but some will get worse despite treatment.

The goals of treatment are to:

- Maintain good organ function
- Lessen symptoms
- Prevent organ damage

Medication

Several medicines are used to treat sarcoidosis.

Corticosteroids: Corticosteroids, which work to reduce inflammation, are the main treatment. Generally, prednisone (a tablet) is given daily or every other day, depending on the symptoms. Prednisone can decrease symptoms, improve lung function, reduce granuloma formation, and possibly lessen scarring of the lungs. Prednisone can be associated with a number of side effects. Because of this, your doctor will carefully monitor you. Prednisone is not the medication of choice for long-term management of sarcoidosis.

Methotrexate: For long-term management of sarcoidosis, steroid-sparing agents are often used. Methotrexate is an anti-inflammatory medicine. It is often used as a second-line medicine. It may be used with corticosteroids or after stopping corticosteroids.

Other Medicines: Other medicines are used if corticosteroids and methotrexate are not effective. These other medicines are not used often, since their effect on sarcoidosis is not as well understood. They also can have side effects. These medicines can include:

- Cellcept®(mycophenolate)
- Humira®(adalimumab)
- Imuran®(azathioprine)
- Plaquenil® (hydroxychloroquine)
- Remicade® (infliximab).

Oxygen Therapy

Oxygen therapy may be an important part of a treatment plan for people with severe sarcoidosis. It can help reduce heart and lung long-term problems caused by low oxygen levels.

Pulmonary Rehabilitation

For people who develop chronic, progressive sarcoidosis, pulmonary rehabilitation also may be helpful. This includes exercise, healthy eating and education.

Because treatment is so important, a person can improve the outcome of sarcoidosis by seeing a doctor when the symptoms first appear. This can help prevent damage to the lungs, eyes, heart and other organs. Also, people with sarcoidosis should continue to follow up with their doctor after they have been diagnosed to monitor if the disease is progressing.

Sarcoidosis at National Jewish Health

National Jewish Health is recognized by the World Association for Sarcoidosis and Other Granulomatous Diseases (WASOG) as a WASOG Sarcoidosis Clinic. This designation provides formal recognition of our team's commitment to meet the needs of sarcoidosis patients and success in keeping to keep abreast of the ongoing advances

and findings.

National Jewish Health is currently involved with ongoing research regarding sarcoidosis. Results of this research will be used to develop better diagnosis and treatment for people with sarcoidosis.

Note: This information is provided to you as an educational service of LUNG LINE® (1-800-222-LUNG). It is not meant to be a substitute for consulting with your own physician.

©Copyright 1989, Revised 1993, 2001, 2005, 2008, 2014, 2018 NATIONAL JEWISH HEALTH. PTE.042