

Pediatric Vaccination Schedule

Vaccines are important to prevent getting serious illnesses. Here is a list of recommended pediatric vaccines from birth to age 18.
Please talk with your child's doctor about the vaccines your child needs and when.

Vaccines	Birth	1 Month	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	19-23 Months	2-3 Years	4-6 Years	7-10 Years	11-12 Years	13-18 Years
DTaP			•	•	•		■				•		•	
HepA						■								
HepB	•	■			■									
Hib			•	•	•	■								
HPV													•	
Influenza					Yearly									
IPV			•	•	■							•		
MMR						•	•				•			
MCV4													•	Booster
PCV			•	•	•	■								
RV			•	•	•									
Varicella						•	•				•			

• Given at these ages

■ The vaccine can be given once during the age range

DTaP (Diphtheria, Tetanus and Pertussis)

HepA (Hepatitis A)

HepB (Hepatitis B)

Hib (*Haemophilus Influenzae* Type b)

HPV (Human Papillomavirus)

Influenza (Flu)

IPV (Polio)

MMR (Measles, Mumps and Rubella)

MCV4 (Meningococcal)

PCV (Pneumococcal)

RV (Rotavirus)

Varicella (Chickenpox)



Sources: The American Academy of Pediatrics, the Centers for Disease Control

5489.10715.MKT8/15

njhealth.org

1.877.CALL NJH (877.225.5654)

© 2015 National Jewish Health

 **National Jewish Health**
Science Transforming Life®