

TESTfacts

An Educational Health Series From National Jewish Health®



REGISTER NOW

to Receive **FREE** Health
Tips, Research Findings,
Ways to Support Our
Mission, News &
More in Your E-Mail Box

Visit njhealth.org/e-news
for more information.

Follow us online!

facebook.com/NJHealthyoutube.com/NationalJewishtwitter.com/njhealth

Two Day Liquid Diet for Constipation

Begin Miralax twice daily, per package directions, 7 days prior to procedure. You will not take it on the day of your prep (day before procedure).

Start a 2 Day Liquid Diet

Start a liquid (no solid foods) diet from the moment of waking up on the day before the procedure.

Liquid Diet – Day 1

- Gatorade, fruit juice (no citrus), soda, popsicles without fruit or cream, gelatin without fruit, broth without fat, coffee, and tea
- No solid food
- No red, orange, or purple colored liquids
- Dairy is ok (as long as it is liquid at room temperature) – including yogurt.

Liquid Diet – Day 2

- Gatorade, fruit juice (no citrus), soda, popsicles without fruit or cream, gelatin without fruit, broth without fat, coffee, and tea
- No solid food
- No red, orange, or purple colored liquids
- No Dairy Allowed

Follow instructions for the prep you were given on Day 2, day before the procedure.

Do not eat or drink (not even water) anything 4 hours prior to your procedure.

- Be sure to inform your doctors and nurses about any diabetic medications and/or blood thinners you take.
- Take all medications for high blood pressure, heart conditions, or seizure disorder with water 4 hours prior to your procedure. This will be the last liquid you should have.
- Bring rescue inhalers, asthma inhalers, diabetic medications, glucose monitor, and CPAP/BiPap machine with you to procedure.
- Please be sure to have a ride arranged and with you to take you home after the procedure. If you do not have a ride with you, your procedure may be canceled.

Call (303)398-1355 option #4, if you have any questions or need to reschedule. ©PTE319, 2/15