

Anticoagulant Medicine Potential for Drug - Food Interactions

Anticoagulant medicines are a group of medicines that inhibit blood clotting, helping to prevent blood clots. Blood clots can cause heart attacks and strokes. Anticoagulant medicines can be used with a number of diseases when there is an increased risk of blood clots. Common anticoagulant medicines include:

- Coumadin®, Jantoven® (warfarin)
- Lovenox® (enoxaparin)
- Hep-Lock U/P®, Hep-Lock®, HepFlush-10® (heparin)



There are important medicine and diet considerations when taking an anticoagulant medicine. There are medications, foods, herbs and vitamins that can interact with anticoagulation medicine. Avoid making any dramatic changes to your diet when taking anticoagulation medicine. We want you to be aware these drug-food interactions when taking anticoagulant medicine. Ask your healthcare provider if you have any questions about these or other possible drug-food interactions. Make sure to discuss any new medication or diet changes with your health care provider before making the change.

Anticoagulants and Medicines

There are many medicines that can interact with anticoagulants. Some medicines can increase the effect of anticoagulants, thus increasing the chances of bleeding. Some medications can decrease the effect of anticoagulants, thus increasing the chances of blood clots.

Antibiotics

Be sure to contact the anticoagulation clinic prior to starting any antibiotic, even if it is for just a few days. Antibiotics can significantly change your INR -- often times very quickly.

Over the counter pain relievers

Over the counter pain relievers that can also increase the effect of anticoagulants, thus increasing the chance of bleeding include:

- Aspirin
- Advil, Motrin, Nuprin (ibuprofen)
- Aleve (naproxen)

Other medications that **increase** the effect of anticoagulants, thus increasing the chance of bleeding include the following:

Allopurinol	Capecitabine	Clopidogrel
Amiodarone	Cephalosporins	Diclofenac
Anabolic steroids	Cimetidine	Disulfiram
Aspirin	Ciprofloxacin	Erythromycin
	Clofibrate	Fluconazole

Medications that **decrease** the effect of anticoagulants, thus increasing the chance of blood clots include the following:

Azathoprine	Griseofulvin	Phenobarbital
Antithyroid medications	Griseofulvin	Rifampin
Carbamazepine	Haloperidol	Vitamin K
Dicloxacillin	Nafcillin	
Glutethimide	Oral contraceptives	

Anticoagulants and Vitamin K

When taking anticoagulants, it is recommended to keep vitamin K intake consistent on a daily basis. Avoid sudden changes in the intake of foods and vitamins containing vitamin K. Consider how much vitamin K you get throughout the day. Read labels of vitamins, minerals and food supplements to identify products high in vitamin K. In general, leafy green vegetables, certain beans and oils are high in vitamin K. Examples of foods and beverages high in vitamin K include:

Beef liver	Cheese	Lentils
Broccoli	Collard greens	Lettuce

Brussel sprouts	Green tea	Spinach
Cabbage	Kale	Soybean oil Turnip greens

Anticoagulants and Alcohol

Food and alcohol may change the way anticoagulants affect your body. Limit the amount of alcohol you drink. If you have a change in your alcohol intake your INR level may need to be checked. Talk with your healthcare provider before you make any changes in the amount of alcohol you drink.

Herbal Supplements and Vitamins

Herbs and vitamins can interact with anticoagulant medication. Remember, consistent use of a multivitamin with low doses of vitamin K. Avoid vitamins that contain high doses of vitamin K and E.

Fish oil may increase the INR level. Discuss the use of fish oil with your health care provider. Your INR level may need to be checked when these changes are made.

Consult your healthcare provider before taking herbal remedies. Herbs can easily interact with your medicines, and if taken together may be harmful to your health. Herbs may also be harmful for certain health conditions or medical procedures. When you see your healthcare provider, inform him/her of which herbs you use.

In summary, there are medications, foods, herbs and vitamins that can interact with anticoagulation medication. Make sure to discuss any new medication, herbal supplements, vitamins or diet changes with your health care provider before making the change.

Visit our website for more information about support groups, clinical trials and lifestyle information.

PTE.196 © Copyright 2010, 2013

NOTE: This information is provided to you as an educational service of National Jewish Health. It is not meant to be a substitute for consulting with your own physician.

National Jewish Health is the leading respiratory hospital in the nation. Founded 119 years ago as a nonprofit hospital, National Jewish Health today is the only facility in the world dedicated exclusively to groundbreaking medical research and treatment of patients with respiratory, cardiac, immune and related disorders. Patients and families come to National Jewish Health from around the world to receive cutting-edge, comprehensive, coordinated care. To learn more, visit njhealth.org.