

Secretin Stimulation Test

What is a secretin stimulation test?

Your doctor has suggested you have a secretin stimulation test as part of the evaluation at National Jewish Health. Secretin is a hormone that is produced by the small intestine when partially digested food is present. When secretin is given to you during this test, certain chemicals may be increased in your blood. This is measured during the test. This test can help your doctor determine a diagnosis and the best treatment for you.

How do you get ready for the test?

- **Do not eat or drink anything after midnight, the night before the test.** No food or drink is allowed until the test is completed.
- Let your doctor know about all the medicine you take.
- The doctor who prescribes your insulin may have you change the dose the day before and the day of the test.
- Let your doctor know if you are taking Atropine as this may decrease the test response.
- Let your doctor know if you are taking Cimetidine, Ranitidine, Catecholamines, Insulin and Haloperidol as they may increase the test response.
- Let your doctor know if you have acute pancreatitis.
- **Acid suppressive medicines are stopped before the test to avoid effects on gastrin release.**
- Stop Proton Pump inhibitors **5 DAYS** before your appointment:
 - Prilosec® (omeprazole), Prevacid® (lansoprazole), Nexium® (esomeprazole), Protonix®, Pantoloc® (pantoprazole), Aciphex® (rabeprazole), Dexilant® (dexlansoprazole).
- Stop H2 Receptor Antagonist **2 DAYS** before your appointment:
 - Zantac® (ranitidine), Tagamet® (cimetidine), Axid® (nizatidine), Pepcid® (famotidine)
- Stop Over the Counter antacids and anti-reflux medications **24 HOURS** before your appointment:
 - Tums, Rolaids, calcium carbonate antacids, Gaviscon, Maalox
- **DO** take any heart, blood pressure or seizure medications **at least 2 hours prior** to leaving home for test with a few small sips of water.
- If you are diabetic, please bring your glucose meter, test strips, and a source of fast-acting glucose with you (such as glucose tablets or glucose gel).

What is done during the test?

When you arrive the nurse will explain what will be done before, during and after the test. If you have any questions, please ask. The nurse will place an IV catheter in your arm. The IV will be used to give you secretin during the test. Blood will be drawn periodically from the IV during the test. Once all the blood samples have been withdrawn, the nurse will remove the IV catheter from your arm and place a dressing on the site. You will be given instructions on signs and symptoms of infection at the IV site before you are discharged.

What should you do after the test?

You can resume normal diet and activities after the test as directed by your doctor.

How long will the test take and when will you find out the results?

The secretin stimulation test takes about 1 hour. You will be notified of your test results about 3 weeks after your test.

How do you get to your test?

If you are being seen at National Jewish Health, on the day of your scheduled test, check in at the front desk. If you have questions or need to cancel or change your appointment, please call (303)-398-1355.

Visit our website for more information about support groups, clinical trials and lifestyle information.

PTE.310 © Copyright 2014 Revised 4/2015

NOTE: This information is provided to you as an educational service of National Jewish Health. It is not meant to be a substitute for consulting with your own physician.

National Jewish Health is the leading respiratory hospital in the nation. Founded 119 years ago as a nonprofit hospital, National Jewish Health today is the only facility in the world dedicated exclusively to groundbreaking medical research and treatment of patients with respiratory, cardiac, immune and related disorders. Patients and families come to National Jewish Health from around the world to receive cutting-edge, comprehensive, coordinated care. To learn more, visit njhealth.org.