T09 10

INHALER MISTAKES

Inhaled asthma medicine needs to reach the airways to work.

Here are 10 common mistakes made when using a
metered-dose inhaler (MDI) and how to correct them.



SITTING DOWN

FIX IT: Standing allows the lungs to fully inhale and provides more power to exhale.





USING AN EMPTY INHALER

FIX IT: Request a refill when the inhaler is half full so you never run out.



NOT SHAKING OR PRIMING THE INHALER

FIX IT: Shake the inhaler canister 10 to 15 times for the medication to be ready to work. When using a new inhaler, prime it by releasing three to four test sprays. Prime again if not used for several weeks.





NOT USING A SPACER WITH AN MDI INHALER

FIX IT: A spacer helps deliver the medication to the airways instead of the mouth. Insert the inhaler into the spacer. Spray one puff of medicine and inhale slowly. Hold your breath for a count of 10 and exhale slowly.



HOLDING THE HEAD TOO FAR FORWARD OR BACKWARD

FIX IT: The head needs to be in a normal position, not too far back or too far forward, to help make a direct path for the medicine to reach the airways.



TONGUE OR TEETH IN THE WAY OF SPACER/INHALER OPENING

FIX IT: Put the spacer/inhaler in the mouth above the tongue, under the top teeth.





M

MOUTH NOT TIGHT ENOUGH AROUND SPACER/INHALER

FIX IT: Close the lips around the spacer so air does not escape.



DIRECTING SPACER/INHALER AT TONGUE OR ROOF OF MOUTH

FIX IT: Aim the spacer/inhaler at the back of the throat, so the medicine reaches the lungs.



SPRAYING SEVERAL PUFFS OF INHALER INTO SPACER

FIX IT: Spray only one puff of the inhaler at a time into the spacer. Breathe out before inhaling. Hold breath for a count of 10, then exhale. Repeat for the number of puffs the doctor prescribed.



10

INHALING MEDICINE TOO FAST

FIX IT: Inhale slowly. A whistle from the spacer means the inhalation is too fast.



REMINDERS

- Follow instructions for using your inhaler.
- Gargle and spit after inhaling a steroid.
- Show your inhaler technique to your provider.
 Clean the inhaler and spacer according.
- Clean the inhaler and spacer according to instructions.

ASTHMA FACTS

- 6.3 million children have asthma.
- Leading chronic disease among children.
- 1 out of 10 school-aged children have asthma.Top cause of missed school days.

